

## Anti-Human ULBP2 Monoclonal Antibody BUMO1

Antigen: Human ULBP2 (UL16-binding protein 2)

Clone: BUMO1, mouse IgG1

**Catalog Number:** BUMO1-500

Specificity: binds: ULBP2

binds not: ULBP1,ULBP3,ULBP4

**Epitope:** in ULBP2 ectodomain

**Applications:** Flow cytometry

Size: 500 μg, 1.0 mg/ml, in 0.5 ml phosphate-buffered saline, pH 7.4 with 0.05%

> sodium azide (Caution: Sodium azide yields highly toxic hydrazoic acid under acidic conditions. Dilute azide compounds in running water before discarding to avoid accumulation of potentially

explosive deposits in plumbing).

In general, for flow cytometry we recommend a final dilution of 10µg mAb/ml **Usage:** 

and for ELISA 1-10 µg mAb/ml.

Purification: Protein A affinity chromatography

Storage: Store at 4°C. For long-term storage freezing at -80°C is recommended.

**Description:** UL16-binding proteins (ULBP) have been discovered in 2001 during a search

for human proteins binding the Human Cytomegalovirus-encoded UL16 glycoprotein [1] and for human homologues of the mouse RAE1 ligands of NKG2D, respectively [2]. ULBP1-4 are cell surface proteins with an MHC class I-like  $\alpha 1/\alpha 2$  superdomain that is bound by human NKG2D [1-3]. ULBP1-3 are attached to the cell surface by GPI-anchor [1]. Expression of ULBP is induced by infection with Human Cytomegalovirus (HCMV) [4]. In vivo expression of ULBP2 is mostly unexplored, except that freshly isolated leukemias have been shown to express ULBP2 [5]. ULBP2 is released from tumor cells by metalloproteases in a manner similar to MIC molecules and can be found in sera of some leukaemia patients [6]. Like other human and mouse NKG2D-

ligands, ULBP stimulate tumor immunity in vivo [7].

Conditions: For research use only. Not for use in diagnostic or therapeutic

procedures. BAMOMAB is not responsible for any patent infringements

caused by the use of this product.

**Country of Origin:** Germany

Literature: 1. Cosman et al. Immunity 14,123-133 (2001).

2. Steinle A et al. Immunogenetics 53, 279-287 (2001).

3. Radaev S et al. Immunity 15,1039-1049 (2001).

4. Welte S et al. Eur J Immunol 33, 194-203 (2003).

5. Salih HR et al. Blood 102, 1389-1396 (2003).

6. Waldhauer I et Steinle A. Cancer Res 66, 2520-2526 (2006).

7. Sutherland C et al. Blood 108:1313-1319 (2006).

Human cell line 293T stained with BUMO1 (black) or IgG1 isotype (dotted) and goat anti-mouse Ig-PE conjugate.